

I searched the following databases using author and keyword suggested by the examiner:

**ProQuest** [Return to the US](#)

Basic **Advanced** Browse My Research [Language](#) [English](#)

Databases selected: Dissertations & Theses

**Results**

6 documents found for: *AkiOkuno or Yabuta or Ooi or Kinoshita* AND *(Omp: variant or mutant or mutated or mutation or mutating or mutagenesis or substitution or substitute or substituted or substituting)* [Refine Search](#) | [Set Up Alert](#) [pg](#)

**Dissertations**


☐ Mark all [Copy](#) [Print](#) [Email](#) [Cite](#) [Export](#) ☐ Show only full text Sort results by: [Mos](#)

- ☐ 1. **Yeast genetics on microarrays**  
by Ooi, Siew Loon, Ph.D., **The Johns Hopkins University**, 2003, 211 pages, AAT 3980738  
[Abstract](#) | [Preview](#) (510 K) | [Full Text](#) : PDF (7 MB) | [Order a copy](#)
- ☐ 2. **Literature of Chinese American and British Chinese writers: immigration policy, citizenship, and racialization**  
by Kinoshita-Baerfort, Anni, Ph.D., **State University of New York at Binghamton**, 2002, 264 pages, AAT 3059844  
[Abstract](#) | [Preview](#) (705 K) | [Full Text](#) : PDF (10 MB) | [Order a copy](#)
- ☐ 3. **The application of production frontier techniques to the metaproduction function: Estimating international agricultural productivity and efficiency**  
by Kinoshita, Randal Mitsugi, Ph.D., **University of Georgia**, 1998, 342 pages, AAT 9828386  
[Abstract](#) | [Preview](#) (543 K) | [Full Text](#) : PDF (13 MB) | [Order a copy](#)
- ☐ 4. **The characterization of protein interaction domains of Goh10, Numb and K11**  
by Ooi, James Yeow-Theng, Ph.D., **New York University**, 1998, 165 pages, AAT 9907185  
[Abstract](#) | [Preview](#) (436 K) | [Full Text](#) : PDF (5 MB) | [Order a copy](#)
- ☐ 5. **AQUEOUS SOLUTION CHEMISTRY OF MOLYBDENUM AND TUNGSTEN CLUSTER COMPLEXES**  
by LEAN, OOI BEE, Ph.D., **University of Newcastle Upon Tyne (United Kingdom)**, 1988, 187 pages, AAT DX84990  
[Abstract](#)
- ☐ 6. **EFFECTS OF CHLORINATED HYDROCARBON INSECTICIDES AND SUBSTITUTED UREA HERBICIDES ON HEPATIC MICROSOOME ENZYMES**  
by KINOSHITA, FLORENCE KEIKO, Ph.D., **The University of Chicago**, 1968, AAT T-17389  
[Citation](#) | [Order a copy](#)

1-6 of 6

**DialogWeb** [Home](#) [About](#) [Contact](#) [Help](#) [Privacy](#) [Terms](#) [Feedback](#) [Search](#)

[Top](#)

 **Dynamic Search: Dissertation Abstracts Online**

[Search Form](#)

Search for  in

Restrict to ☒ Masters Theses  
☒ Doctoral Dissertations

Published from  To  (YYYY)

Browse List of

► No Records Found : Dissertation Abstracts Online

 **warning**

There were no records matching your search. Please modify your search and try again.

#### Tips:

- Don't over-specify: use only the search options you really need.
- Exclude "implied concepts": leave out words like *research* or *effects*.
- Check the format of your entry: some search options require specific spacing or punctuation.
- Use the Browse feature when available to find variations for your terms.
- Use more wildcards to search different word endings.
- Check that you are using parentheses correctly when you combine words with AND, OR, NOT.
- Check for misspelled words.

[refine search](#)



Searching: Academic Search Premier, Show all | [Choose Databases](#)

Okuno or Yabuta or Ooi or Kinoshita in AU Author

[Search](#) [Clear](#)

Ompt variant or mutant or mutated or mutation or mutating or mutagenesis or substitution or substitute or substituted or substituting

AND in TI Title


AND in Select a field (optional) [Add Row](#)

[Basic Search](#) | [Advanced Search](#) | [Visual Search](#) | [Search History](#)


**10 Results for...**

**Boolean/Phrases:**  
AU ( Okuno or Yabuta or Ooi or Kinoshita ) and TI ( Ompt variant ...

**Expanders**

 Also search within the full text of the articles

**Limiters**


 Published Date from: 2003/01/01-2014/12/31


**1. Utilization of *Escherichia coli* Outer-Membrane Endoprotease OmpT Variants as Processing Enzymes for Production of Peptides from Designer Fusion Proteins.** [Add](#)

By: **Okuno, Kazuko; Yabuta, Masayuki; Ooi, Toshihiro; Kinoshita, Shinichi**, *Applied & Environmental Microbiology*, Jan2014, Vol. 70 Issue 1, p76-86, 11p, 9 Diagrams, 2 Charts, 1 Graph; DOI: 10.1128/AEM.70.1.76-86.2014


Subjects: *ESCHERICHIA coli*; ENZYMES; PEPTIDES; PROTEINS; AMINO acids; PROTEOLYTIC enzymes

Database: Academic Search Premier

 [Add to Folder](#) | Relevancy: **xxxxxxxx**

 [Linked Full Text](#)

[PubMed](#)
[Resources](#)
[How To](#)


 U.S. National Library of Medicine  
 National Institutes of Health

Search: PubMed

[RSS](#)
[Save search](#)
[Limits](#)
[Advanced search](#)
[Help](#)

Display Settings: Abstract

Send to:

We found 1 article by title matching your search:

[Asai Environ Microbiol. 2004 Jan;70\(1\):78-86.](#)

# **Utilization of Escherichia coli outer-membrane endoprotease OmpT variants as processing enzymes for production of peptides from designer fusion proteins.**

[Okuno K](#), [Yabuta M](#), [Ooi T](#), [Kinoshita S](#)

Institute for Medicinal Research and Development, Daiichi Sankyo Pharma Co., Ltd., Akaawa, Chiyoda-machi, Ohta-gun, Gunma 370-0503, Japan.  
[Kazuaki.Okuno@dsup.co.jp](#)

## **Abstract**

Escherichia coli outer-membrane endoprotease OmpT has suitable properties for processing fusion proteins to produce peptides and proteins. However, utilization of this protease for such production has been restricted due to its generally low cleavage efficiency at Arg (or Lys)-Xaa, where Xaa is a nonbasic N-terminal amino acid of a target polypeptide. The objective of this study was to generate a specific and efficient OmpT protease and to utilize it as a processing enzyme for producing various peptides and proteins by converting its substrate specificity. Since OmpT Asp(97) is proposed to interact with the P1' amino acid of its substrates, OmpT variants with variations at Asp(97) were constructed by replacing this amino acid with 19 natural amino acids to alter the cleavage specificity at Arg (P1)-Xaa (P1'). The variant OmpT that had a methionine at this position, but not the wild-type OmpT, efficiently cleaved a fusion protein containing the amino acid sequence -Arg-Arg-Ala-Arg downward arrow motilin, in which motilin is a model peptide with a phenylalanine at the N terminus. The OmpT variants with leucine and histidine at position 97 were useful in releasing human adrenocorticotrophic hormone (1-24) (serine at the N terminus) and human calcitonin precursor (cysteine at the N terminus), respectively, from fusion proteins. Motilin was produced by this method and was purified up to 99.0% by two chromatographic steps; the yield was 160 mg/liter of culture. Our novel method in which the OmpT variants are used could be employed for production of various peptides and proteins.

PMID- 14711628

OWN - NLM

STAT- MEDLINE

DA - 20040108

DCOM- 20040408

LR - 20100617

IS - 0099-2240 (Print)

IS - 0099-2240 (Linking)

VI - 70

IP - 1

DP - 2004 Jan

TI - Utilization of Escherichia coli outer-membrane endoprotease OmpT variants as

processing enzymes for production of peptides from designer fusion proteins.

PG - 76-86

AB - Escherichia coli outer-membrane endoprotease OmpT has suitable properties for

processing fusion proteins to produce peptides and proteins.

However, utilization

of this protease for such production has been restricted due to its generally low

cleavage efficiency at Arg (or Lys)-Xaa, where Xaa is a nonbasic N-terminal amino

acid of a target polypeptide. The objective of this study was to generate a

specific and efficient OmpT protease and to utilize it as a processing enzyme for

producing various peptides and proteins by converting its substrate specificity.

Since OmpT Asp(97) is proposed to interact with the P1' amino acid of its substrates, OmpT variants with variations at Asp(97) were constructed by replacing this amino acid with 19 natural amino acids to alter the cleavage specificity at Arg (P1)-Xaa (P1'). The variant OmpT that had a methionine at this position, but not the wild-type OmpT, efficiently cleaved a fusion protein containing the amino acid sequence -Arg-Arg-Arg-Ala-Arg downward arrow motilin, in which motilin is a model peptide with a phenylalanine at the N terminus. The OmpT variants with leucine and histidine at position 97 were useful in releasing human adrenocorticotrophic hormone (1-24) (serine at the N terminus) and human calcitonin precursor (cysteine at the N terminus), respectively, from fusion proteins. Motilin was produced by this method and was purified up to 99.0% by two chromatographic steps; the yield was 160 mg/liter of culture. Our novel method in which the OmpT variants are used could be employed for production of various peptides and proteins.

AD - Institute for Medicinal Research and Development, Daiichi Suntory Pharma Co.,

Ltd., Akaiwa, Chiyoda-machi, Ohra-gun, Gunma 370-0503, Japan.

Kazuaki\_Okuno@dsup.co.jp

FAU - Okuno, Kazuaki

AU - Okuno K

FAU - Yabuta, Masayuki

AU - Yabuta M

FAU - Ooi, Toshihiko

AU - Ooi T

FAU - Kinoshita, Shinichi

AU - Kinoshita S

LA - eng

PT - Journal Article

PL - United States

TA - Appl Environ Microbiol

JT - Applied and environmental microbiology

JID - 7605801

RN - 0 (Bacterial Outer Membrane Proteins)

RN - 0 (Escherichia coli Proteins)

RN - 0 (Peptides)

RN - 0 (Porins)

RN - 0 (Recombinant Fusion Proteins)

RN - 0 (ompT protein, E coli)

RN - 52906-92-0 (Motilin)

RN - EC 3.4.- (Peptide Hydrolases)

RN - EC 3.4.21.- (Serine Endopeptidases)

SB - IM

MH - Amino Acid Sequence

MH - Bacterial Outer Membrane Proteins

MH - Escherichia coli/\*enzymology/genetics  
 MH - Escherichia coli Proteins  
 MH - \*Genetic Variation  
 MH - Humans  
 MH - Inclusion Bodies/metabolism  
 MH - Molecular Sequence Data  
 MH - Motilin/genetics/metabolism  
 MH - Mutation  
 MH - Peptide Hydrolases  
 MH - Peptides/\*metabolism  
 MH - Porins/genetics/\*metabolism  
 MH - Recombinant Fusion Proteins  
 MH - Serine Endopeptidases/genetics/\*metabolism  
 MH - Substrate Specificity  
 PMC - PMC321264  
 OID - NLM: PMC321264  
 EDAT- 2004/01/09 05:00  
 MHDA- 2004/04/09 05:00  
 CRDT- 2004/01/09 05:00  
 PST - ppublish  
 SO - Appl Environ Microbiol. 2004 Jan;70(1):76-86.

**scirus**  
 for scientific information only

Advanced search

Any of the words  in

AND

Any of the words  in

**scirus**  
 for scientific information only

Advanced

1-15 of 753 hits for {author:Okuno OR author:Yabuta OR author:Ooi OR author:Kinoshita} {title:Omp OR title:t OR title:variant OR title:mutant...

or  checked results

**Filter search results by**

**Content sources**

☒ **Journal sources (669)**

- MEDLINE / PubMed (424)
- ScienceDirect (141)
- Pubmed Central (34)

[more >](#)

☒ **Preferred web (182)**

- Patent Offices (78)
- Digital Archives (20)
- The University of Hong Kong (5)

[more >](#)


☐ **Other web (12)**

**File types**

- HTML (782)
- PDF (154)


Did you mean {author:Okuno OR author:abuta OR author:Ooi OR author:kinoshita} {title:Omp OR title:t OR title:variant OR title:mutant OR title:mutated OR title:mutation OR title:mutating OR title:mutagenesis OR title:substitution OR title:substitute OR title:substituted OR title:substituted}?

- ☒ 1. BCMA modulates T cell responses and is involved in autoimmune disease  
 Muramatsu, Rieko / Kubo, Takekazu / Mori, Masahiro / Nakamura, Yuka / Fujita, Yuki / Akutsu, Tsugio / Okuno, Tatsusada / [...] / Yamashita, Toshihide, *Nature Medicine*, 17 (4), p.486-494, Apr 2011  
 doi:10.1038/nm.2321

Published journal article available from  nature publishing group

similar results

- ☒ 2. Ataxia-telangiectasia mutated (ATM) deficiency decreases reprogramming efficiency and leads to genomic instability in iPS cells  
 Kinoshita, Taisuke / Nagamatsu, Go / Kosaka, Takeo / Takubo, Kaiyo / Hotta, Akitsu / Ellis, James / Suda, Toshio, *Biochemical and Biophysical Research Communications*, 407 (2), p.321, Apr 2011  
 doi:10.1016/j.bbrc.2011.03.013  
 Abstract During cell division, one of the major features of somatic cell reprogramming by defined factors, cells are potentially exposed to DNA damage. Inactivation of the tumor suppressor gene p53 raised reprogramming efficiency but resulted in ...

Published journal article available from  ScienceDirect

Same database using AND for authors and no words in the title section:

**Scirus**  
for scientific information only

author:Okuno AND author:Yabuta AND author:Ooi AND author:Kinoshita

1-5 of 5 hits for author:Okuno AND author:Yabuta AND author:Ooi AND author:Kinoshita

Filter search results by

Content sources

Journal sources (4)

MEDLINE / PubMed (3)

PubMed Central (1)

Other web (1)

File types

HTML (5)

Refine your search

fusion proteins

cleavage

peptides


Did you mean author:Okuno author:yabuta author:ooi author:kinoshita?

1. Utilization of *Escherichia coli* Outer-Membrane Endoprotease OmpT Variants as Processing Enzymes for Production of Peptides from Designer Fusion Proteins -- Okuno et al. 70 (1): 76 -- Applied and Environmental Microbiology (117K)  
Kazuaki Okuno / Masayuki Yabuta / Toshihiko Ooi / Shinichi Kinoshita . Jun 2004  
Applied and Environmental Microbiology, January 2004, p. 76-86, Vol. 70, No. 1 0099-2240/04/\$08.00+0 DOI: 10.1128/AEM.70.1.76-86.2004 Copyright © 2004, American Society for Microbiology. All Rights Reserved. [http://aem.asm.org/cgi/content/full/70/1/76]  
similar results

2. Utilization of *Escherichia coli* outer-membrane endoprotease OmpT variants as processing enzymes for production of peptides from designer fusion proteins.  
Okuno, Kazuaki / Yabuta, Masayuki / Ooi, Toshihiko / Kinoshita, Shinichi. Applied and environmental microbiology, 70 (1), p.76-86, Jan 2004  
Escherichia coli outer-membrane endoprotease OmpT has suitable properties for processing fusion proteins to produce peptides and proteins. However, utilization of this protease for such production has been restricted due to its generally low cleavage ...

MEDLINE/PubMed Citation on PubMed

The information from the Scirus database first link is below. The second link is to PubMed which is already in this document.

 Applied and Environmental Microbiology

HOME | CURRENT ISSUE | ARCHIVE | ALERTS | ABOUT ASM | CONTACT US | TECH SUPPORT | Journals.ASM.org

Institution: US PATENT & TRADEMARK OFFICE

« Previous Article | Next Article »

Applied and Environmental Microbiology, January 2004, p. 76-86, Vol. 70, No. 1  
0099-2240/04/\$08.00+0 DOI: 10.1128/AEM.70.1.76-86.2004  
Copyright © 2004, American Society for Microbiology. All Rights Reserved.

**Utilization of *Escherichia coli* Outer-Membrane Endoprotease OmpT Variants as Processing Enzymes for Production of Peptides from Designer Fusion Proteins**

Kazuaki Okuno,<sup>1,2\*</sup> Masayuki Yabuta,<sup>1</sup> Toshihiko Ooi,<sup>2</sup> and Shinichi Kinoshita<sup>2</sup>

Institute for Medicinal Research and Development, Daiichi Suntory Pharma Co., Ltd., Akaiwa, Chiyoda-machi, Ohra-gun, Gunma 370-0503,<sup>1</sup> Division of Molecular Chemistry, Graduate School of Engineering, Hokkaido University, Kita-ku, Sapporo, Hokkaido 060-8628, Japan<sup>2</sup>

Received 18 June 2003/ Accepted 13 October 2003

From the Scirus second link above:

